

The Claims:

1. (Previously Presented) An apparatus comprising:
one or more processors; and
a memory coupled to the processors comprising one or more instructions, the processors operable when executing the instructions to:
determine an atom-pair type of a protein-ligand atom pair in a protein-ligand complex;
calculate a repulsion term of the protein-ligand atom pair according to a minimum binding-energy distance value and a well-depth value of the atom-pair type;
calculate a potential of mean force (PMF) of the protein-ligand atom pair according to the calculated repulsion term of the protein-ligand atom pair; and
calculate a PMF score of the protein-ligand complex according to the calculated PMF of the protein-ligand atom pair, the PMF score indicating a binding affinity between a protein and a ligand in the protein-ligand complex.
2. (Currently Amended) The ~~system~~ apparatus of Claim 1, wherein the minimum binding-energy distance value of the atom-pair type is an empirically derived minimum binding-energy distance value of the atom-pair type and the well-depth value of the atom-pair type is an empirically derived well-depth value of the atom-pair type.
3. (Canceled)
4. (Canceled)
5. (Currently Amended) The ~~system~~ apparatus of Claim 2, wherein a first set of empirically derived minimum binding-energy distances and well-depth values comprises the minimum binding-energy distance value and the well-depth value of the atom-pair type, the first set yielding a better agreement with a plurality of actual analyzed protein-ligand atom pairs than one or more second sets of empirically derived minimum binding-energy distances and well-depth values.
6. (Canceled)

7. (Currently Amended) The ~~system~~ apparatus of Claim 5, wherein root mean square (RMS) deviation between actual analyzed protein-ligand complex structures and protein-ligand complex structures predicted according to a set of empirically derived minimum binding-energy distances and well-depth values determines agreement between the set of empirically derived minimum binding-energy distances and well-depth values and the actual analyzed protein-ligand atom pairs.

8. (Canceled)

9. (Currently Amended) The ~~system~~ apparatus of Claim 5, wherein one or more of the first set of empirically derived minimum binding-energy distances and well-depth values or second sets of empirically derived minimum binding-energy distances and well-depth values are each a product of one or more manual processes or automatic processes.

10. (Currently Amended) The ~~system~~ apparatus of Claim 9, wherein at least one of the automatic processes comprises execution of a genetic algorithm.

11. (Previously Presented) A method comprising:
determining an atom-pair type of a protein-ligand atom pair in a protein-ligand complex;

calculating a repulsion term of the protein-ligand atom pair according to a minimum binding-energy distance value and a well-depth value of the atom-pair type;

calculating a potential of mean force (PMF) of the protein-ligand atom pair according to the calculated repulsion term of the protein-ligand atom pair; and

calculating a PMF score of the protein-ligand complex according to the calculated PMF of the protein-ligand atom pair, the PMF score indicating a binding affinity between a protein and a ligand in the protein-ligand complex.

12. (Previously Presented) The method of Claim 11, wherein the minimum binding-energy distance value of the atom-pair type is an empirically derived minimum binding-energy distance value of the atom-pair type and the well-depth value of the atom-pair type is an empirically derived well-depth value of the atom-pair type.

13. (Canceled)

14. (Canceled)

15. (Previously Presented) The method of Claim 12, wherein a first set of empirically derived minimum binding-energy distances and well-depth values comprises the minimum binding-energy distance value and the well-depth value of the atom-pair type, the first set yielding a better agreement with a plurality of actual analyzed protein-ligand atom pairs than one or more second sets of empirically derived minimum binding-energy distances and well-depth values.

16. (Canceled)

17. (Previously Presented) The method of Claim 15, wherein root mean square (RMS) deviation between actual analyzed protein-ligand complex structures and protein-ligand complex structures predicted according to a set of empirically derived minimum binding-energy distances and well-depth values determines agreement between the set of empirically derived minimum binding-energy distances and well-depth values and the actual analyzed protein-ligand atom pairs.

18. (Canceled)

19. (Previously Presented) The method of Claim 5, wherein one or more of the first set of empirically derived minimum binding-energy distances and well-depth values or second sets of empirically derived minimum binding-energy distances and well-depth values are each a product of one or more manual processes or automatic processes.

20. (Previously Presented) The method of Claim 19, wherein at least one of the automatic processes comprises execution of a genetic algorithm.

21. (Previously Presented) Logic encoded in one or more media for execution and when executed operable to:

determine an atom-pair type of a protein-ligand atom pair in a protein-ligand complex;

calculate a repulsion term of the protein-ligand atom pair according to a minimum binding-energy distance value and a well-depth value of the atom-pair type;

calculate a potential of mean force (PMF) of the protein-ligand atom pair according to the calculated repulsion term of the protein-ligand atom pair; and

calculate a PMF score of the protein-ligand complex according to the calculated PMF of the protein-ligand atom pair, the PMF score indicating a binding affinity between a protein and a ligand in the protein-ligand complex.

22. (Previously Presented) The logic of Claim 21, wherein the minimum binding-energy distance value of the atom-pair type is an empirically derived minimum binding-energy distance value of the atom-pair type and the well-depth value of the atom-pair type is an empirically derived well-depth value of the atom-pair type.

23. (Canceled)

24. (Canceled)

25. (Previously Presented) The logic of Claim 22, wherein a first set of empirically derived minimum binding-energy distances and well-depth values comprises the minimum binding-energy distance value and the well-depth value of the atom-pair type, the first set yielding a better agreement with a plurality of actual analyzed protein-ligand atom pairs than one or more second sets of empirically derived minimum binding-energy distances and well-depth values.

26. (Canceled)

27. (Previously Presented) The logic of Claim 25, wherein root mean square (RMS) deviation between actual analyzed protein-ligand complex structures and protein-ligand complex structures predicted according to a set of empirically derived minimum binding-energy distances and well-depth values determines agreement between the set of empirically derived minimum binding-energy distances and well-depth values and the actual analyzed protein-ligand atom pairs.

28. (Canceled)

29. (Previously Presented) The logic of Claim 25, wherein one or more of the first set of empirically derived minimum binding-energy distances and well-depth values or second sets of empirically derived minimum binding-energy distances and well-depth values are each a product of one or more manual processes or automatic processes.

30. (Previously Presented) The logic of Claim 29, wherein at least one of the automatic processes comprises execution of a genetic algorithm.

31. (Previously Presented) A system comprising:
means for determining an atom-pair type of a protein-ligand atom pair in a protein-ligand complex;
means for calculating a repulsion term of the protein-ligand atom pair according to a minimum binding-energy distance value and a well-depth value of the atom-pair type;
means for calculating a potential of mean force (PMF) of the protein-ligand atom pair according to the calculated repulsion term of the protein-ligand atom pair; and
means for calculating a PMF score of the protein-ligand complex according to the calculated PMF of the protein-ligand atom pair, the PMF score indicating a binding affinity between a protein and a ligand in the protein-ligand complex.